arXiv:math/0608425v1 [math.RA] 16 Aug 2006

Some remarks on localization in coalgebras

Gabriel Navarro*

Department of Algebra, Faculty of Science, University of Granada
c. Fuentenueva s/n, 18071 Granada
Spain

Abstract

We analyze the geometry of the Ext-quiver of a coalgebra C in order to study the behavior of simple and injective C-comodules under the action of the functors associated to a localizing subcategory of the category of C-comodules.

Introduction

In recent years, the difficulty of studying coalgebras in a general framework has caused the appearance of several kinds of coalgebras which are investigated separately. Many of them are defined through certain properties involving their category of comodules or, merely, their simple or injective comodules. For instance, this is the case of cosemisimple, pure semisimple, semiperfect, quasi-co-Frobenius, hereditary or serial coalgebras, see [3], [4], [7], [12] or [15]. Therefore it is natural to ask about the behavior of such comodules in different situations. In this paper we shall focus on this question in the context of localization as defined by Gabriel in [5]. In particular, we will study the behavior of simple and (indecomposable) injective comodules under the action of the localizing functors associated to a localizing subcategory. In many cases this analysis is related to the geometry of the Ext-quiver associated to the coalgebra. For that reason, in Section 2 we take into consideration some geometric properties of it. We devote Section 3, 4 and 5 to study how the quotient, section and colocalizing functors

---

Keywords and phrases: coalgebra, localization, Ext-quiver.
2000 *Mathematics Subject Classification*: 18E35, 16W30.
E-mail: gnavarro@ugr.es
*Research supported by Spanish MEC grant BES-2002-2403, DGES BMF2001-2823 and FQM-266 (Junta de Andalucía Research Group).

transform simple and injective objects, respectively. Lastly, in Section 6, we relate those properties to the idempotent associated to the localizing subcategory, extending the results of [10].

1 Preliminaries and notation

Throughout we fix a ground field K and we assume that all vector spaces are over K and every map is a K-linear map. In particular, C is a K-coalgebra which, without loss of generality, we shall consider basic. The reader is expected to be familiar with basic coalgebra theory, see for instance [1], [13] or [19]. Unless otherwise stated, all C-comodules will be right C-comodules. We denote by $\mathcal{M}^C$ the category of right C-comodules and by $\mathcal{M}^C_{qf}$ and $\mathcal{M}^C_f$ the full subcategories of $\mathcal{M}^C$ formed by quasi-finite and finite dimensional comodules, respectively. We also denote by $\{S_x\}_{x\in I_C}$ and by $\{E_x\}_{x\in I_C}$ a complete set of pairwise non-isomorphic simple and indecomposable injective right C-comodules, respectively. For each right C-comodule M, we may calculate its socle, $\operatorname{Soc} M$, and its injective envelope, $E(M)$. Then, for each $x \in I_C$, we assume that $\operatorname{Soc} E_x = S_x$, and consequently, $E(S_x) = E_x$.

We will follow the quiver notation of [8] and [10], in particular, by a quiver we mean an oriented graph $Q = (Q_0, Q_1)$, where Q_0 is set of vertices and Q_1 is the set of arrows. The path coalgebra KQ is defined in the standard way. The set of simple right KQ-comodules is $\{Kx\}_{x\in Q_0}$, and, for each $x \in Q_0$, $E_x = E(Kx)$ is generated by the set of paths ending at x.

Following [5], a dense subcategory $\mathcal{T}$ of $\mathcal{M}^C$ is said to be localizing if the *quotient functor* $T : \mathcal{M}^C \to \mathcal{M}^C/\mathcal{T}$ has a right adjoint functor, S, called the *section functor*. T is exact, S is fully faithful and left exact, and $TS = 1_{\mathcal{M}^C/\mathcal{T}}$. Dually, see [16], $\mathcal{T}$ is said to be colocalizing if T has a left adjoint functor, H, called the *colocalizing functor*. H is a fully faithful and right exact functor such that $TH = 1_{\mathcal{M}^C/\mathcal{T}}$. It is well-known that there exist one-to-one correspondences between localizing subcategories, indecomposable injective comodules and simple comodules. In [2], [10] and [21] localizing subcategories are described by means of idempotent elements of the dual algebra C^*. In particular, it is proved that the quotient category is the category of right comodules over the coalgebra eCe, where e is the idempotent element associated to the localizing subcategory. See the above references for the coalgebra structure of eCe and a description of the localization functors by means of e. The localization in categories of comodules over path coalgebras is described in detail in [10].

2 The geometry of the Ext-quiver

To any coalgebra C, we may associate a quiver Γ_C known as the right Ext-quiver of C, see [14]. We recall that the set of vertices of Γ_C is the set of pairwise non-isomorphic simple right C-comodules $\{S_x\}_{x\in I_C}$ and, for two vertices S_x and S_y, there exists a unique arrow $S_y \to S_x$ in Γ_C if and only if $\mathrm{Ext}^1_C(S_y,S_x)\neq 0$. We may proceed analogously with left C-comodules and obtain the left Ext-quiver of C, Δ_C. Throughout we assume that Γ_C is connected, i.e., C is indecomposable as coalgebra.

Remark 2.1. *The results obtained in this paper are also valid for the valued Gabriel quiver of* C, (Q^C,d^C), *i. e., following [11], the valued quiver whose set of vertices is* $\{S_x\}_{x\in I_C}$ *and such that there exists a unique valued arrow* $S_y \xrightarrow{(d^1_{yx},d^2_{yx})} S_x$ *if and only if* $\mathrm{Ext}^1_C(S_y,S_x)\neq 0$ *and* $d^1_{yx}=\dim_{\mathrm{End}_C(S_y)}\mathrm{Ext}^1_C(S_y,S_x)$ *as right* $\mathrm{End}_C(S_y)$*-module and* $d^2_{yx}=\dim_{\mathrm{End}_C(S_x)}\mathrm{Ext}^1_C(S_y,S_x)$ *as left* $\mathrm{End}_C(S_x)$*-module.*

This is also true if we consider the (non-valued) Gabriel quiver. That quiver is obtained taking the same set of vertices and the number of arrows from a vertex S_x *to a vertex* S_y *is* $\dim_{\mathrm{End}_C(S_x)}\mathrm{Ext}^1_C(S_x,S_y)$. *We recall that if* C *is pointed (or* K *is algebraically closed) then it is isomorphic to a subcoalgebra of the path coalgebra of its Gabriel quiver.*

Let us take into consideration some geometric properties of Γ_C. Given a vertex S_x, we say that the vertex S_y is an *immediate predecessor* (or a 1-predecessor) of S_x if there exists an arrow $S_y \to S_x$ in Γ_C.

Lemma 2.2. S_y *is an immediate predecessor of* S_x *if and only if* $S_y \subseteq \mathrm{Soc}(E_x/S_x)$.

Proof. Let us consider the short exact sequence $S_x \hookrightarrow E_x \to E_x/S_x$. We apply to it the functor $\mathrm{Hom}_C(S_y,-)$ and then we obtain the long exact sequence

$$\begin{tikzcd}
0 \arrow[r] & \mathrm{Hom}_C(S_y,S_x) \arrow[r, "f"] & \mathrm{Hom}_C(S_y,E_x) \arrow[r] & {} \\
{} \arrow[r] & \mathrm{Hom}_C(S_y,E_x/S_x) \arrow[r] & \mathrm{Ext}^1_C(S_y,S_x) \arrow[r] & 0
\end{tikzcd}$$

Since f is a K-linear isomorphism, $\mathrm{Hom}_C(S_y,E_x/S_x)\cong \mathrm{Ext}^1_C(S_y,S_x)$ and the result follows. □

We may generalize the former definition by means of the socle filtration. Following [6], any right C-comodule M has a filtration

$$0 \subset \mathrm{Soc}\, M \subset \mathrm{Soc}^2 M \subset \cdots \subset M$$

called the *Loewy series*, where, for $n > 1$, $\mathrm{Soc}^n M$ is the unique subcomodule of M satisfying that $\mathrm{Soc}^{n-1} M \subset \mathrm{Soc}^n M$ and

$$\frac{\mathrm{Soc}^n M}{\mathrm{Soc}^{n-1} M} = \mathrm{Soc}\left(\frac{M}{\mathrm{Soc}^{n-1} M}\right).$$

Lemma 2.3. *[6] Let M and N be right C-comodules then:*

(a) *Let n be a positive integer. If $0 \subseteq R_1 \subseteq R_2 \subseteq \cdots \subseteq R_n$ is a chain of subcomodules of M such that R_i/R_{i-1} is semisimple for all $i = 1, \ldots, n$ then $R_n \subseteq \mathrm{Soc}^n M$.*

(b) *If N is a subcomodule of M then $\mathrm{Soc}^n N = \mathrm{Soc}^n M \cap N$ for all $n \geq 1$.*

(c) *If $f : N \to M$ is a morphism of right C-comodules then $f(\mathrm{Soc}^n N) \subseteq \mathrm{Soc}^n M$ for all $n \geq 1$.*

(d) *If $M = \bigoplus_\lambda M_\lambda$ then $\mathrm{Soc}^n M = \bigoplus_\lambda \mathrm{Soc}^n M_\lambda$ for all $n \geq 1$.*

We shall need the following result:

Lemma 2.4. *Let M be a right C-comodule. Then, for each positive integer n, the chain*

$$0 \subset \frac{\mathrm{Soc}^{n+1} M}{\mathrm{Soc}^n M} \subset \frac{\mathrm{Soc}^{n+2} M}{\mathrm{Soc}^n M} \subset \cdots \subset \frac{\mathrm{Soc}^{n+t} M}{\mathrm{Soc}^n M} \subset \cdots$$

is the Loewy series of the right C-comodule $M/\mathrm{Soc}^n M$, that is,

$$\mathrm{Soc}^t\left(\frac{M}{\mathrm{Soc}^n M}\right) = \frac{\mathrm{Soc}^{n+t} M}{\mathrm{Soc}^n M} \qquad \text{for each } t > 0$$

Proof. For each $t > 0$, denote by N_t the C-comodule $\mathrm{Soc}^{n+t} M/\mathrm{Soc}^n M$ and by N the C-comodule $M/\mathrm{Soc}^n M$. The case $t = 1$ follows from the definition of the Loewy series. Assume now that the statement holds for $t - 1$. Then N_t is a subcomodule of N such that $\mathrm{Soc}^{t-1} N \subset N_t$ and

$$\frac{N_t}{\mathrm{Soc}^{t-1} N} \cong \frac{\mathrm{Soc}^{n+t} M}{\mathrm{Soc}^{n+t-1} M} \cong \mathrm{Soc}\left(\frac{M}{\mathrm{Soc}^{n+t-1} M}\right) \cong \mathrm{Soc}\left(\frac{N}{\mathrm{Soc}^{t-1} N}\right).$$

Thus $\mathrm{Soc}^t N = N_t$. □

Given a vertex S_x, we say that the vertex S_y is an *n-predecessor* of S_x if $\mathrm{Ext}^1_C(S_y, \mathrm{Soc}^n E_x) \neq 0$, or equivalently, proceeding as in Lemma 2.2, if $S_y \subseteq \mathrm{Soc}\,(E_x/\mathrm{Soc}^n E_x)$.

Convention 2.5. *Throughout, given a simple comodule S_x, for each $n \geq 1$, we shall denote by $\{S_i\}_{i \in I_n}$ the set of simple C-comodules such that*

$$\mathrm{Soc}\left(\frac{E_x}{\mathrm{Soc}^n E_x}\right) = \bigoplus_{i \in I_n} S_i,$$

and we will refer to it as the set of all n-predecessors of S_x. Obviously, each n-predecessor S_i is repeated r_i times, where

$$r_i = \frac{\dim_K \mathrm{Hom}_C(S_i, E_x/\mathrm{Soc}^n E_x)}{\dim_K \mathrm{End}_C(S_i)} = \frac{\dim_K \mathrm{Ext}^1_C(S_i, \mathrm{Soc}^n E_x)}{\dim_K \mathrm{End}_C(S_i)}.$$

Lemma 2.6. *Let S_x and S_y be two simple C-comodules. The following assertions are equivalent:*

- (a) *S_y is a n-predecessor of S_x.*
- (b) *There exists a non-zero morphism $f : \mathrm{Soc}^{n+1} E_x \to E_y$ such that $f(\mathrm{Soc}^n E_x) = 0$.*
- (c) *There exists a morphism $g : E_x \to E_y$ such that $g(\mathrm{Soc}^i E_x) = 0$ for all $i = 1, \ldots, n$ and $g(\mathrm{Soc}^{n+1} E_x) \neq 0$*

Proof. $(a) \Leftrightarrow (b)$. Assume that S_y is an n-predecessor of S_x. Then there exists a non-zero map $h : \mathrm{Soc}^{n+1} E_x/\mathrm{Soc}^n E_x \to E_y$ making commutative the following diagram

$$\begin{tikzcd} S_y \arrow[r, hook, "i"] \arrow[d, hook, "i"'] & \frac{\mathrm{Soc}^{n+1} E_x}{\mathrm{Soc}^n E_x} \arrow[dl, dashed, "h"] \\ E_y & \end{tikzcd}$$

Then, the composition $\mathrm{Soc}^{n+1} E_x \xrightarrow{p} \mathrm{Soc}^{n+1} E_x/\mathrm{Soc}^n E_x \xrightarrow{h} E_y$ is a nonzero morphism which vanishes in $\mathrm{Soc}^n E_x$.

Conversely, given such an f, it decomposes through a non-zero morphism $g : \bigoplus_{i \in I_n} S_i \longrightarrow E_y$. Therefore there is an $i \in I_n$ such that $gu_i : S_i \to E_y$ is non-zero, where u_i is the standard inclusion. That is, $S_i = S_y$ is an n-predecessor of S_x.

$(b) \Leftrightarrow (c)$. Assume that f is such a morphism. Since E_y is an injective C-comodule, there exists a morphism $g : E_x \to E_y$ such that

$$\begin{tikzcd} \mathrm{Soc}^{n+1} E_x \arrow[r, hook, "i"] \arrow[d, "f"'] & E_x \arrow[dl, dashed, "g"] \\ E_y & \end{tikzcd}$$

commutes. Obviously, $g(\mathrm{Soc}^n E_x) = 0$ and $g(\mathrm{Soc}^{n+1} E_x) \neq 0$.

For the converse, it is enough to consider the restriction of g to the subcomodule $\mathrm{Soc}^{n+1} E_x$. □

Remark 2.7. *Observe that the proof is also valid for an arbitrary C-comodule M, that is, the following assertions are equivalent:*

(*a*) $S_y \subseteq M/\mathrm{Soc}^n M$.

(*b*) *There exists a non-zero morphism* $f : \mathrm{Soc}^{n+1}M \to E_y$ *such that* $f(\mathrm{Soc}^n M) = 0$.

(*c*) *There exists a morphism* $g : M \to E_y$ *such that* $g(\mathrm{Soc}^i M) = 0$ *for all* $i = 1, \ldots, n$ *and* $g(\mathrm{Soc}^{n+1}M) \neq 0$

We simply say that S_y is a predecessor of S_x if there exists an integer $n \geq 1$ such that S_y is an n-predecessor of S_x. The following result gives a necessary and sufficient condition for the vertex S_y to be a predecessor of a vertex S_x. Given two indecomposable injective right C-comodules E_x and E_y, we denote by $\mathrm{Rad}_C(E_x, E_y)$ the set of all morphisms in $\mathrm{Hom}_C(E_x, E_y)$ which are not bijective.

Corollary 2.8. *Let S_x and S_y be two simple C-comodules. Then, S_y is a predecessor of S_x if and only* $\mathrm{Rad}_C(E_x, E_y) \neq 0$.

Proof. The sufficiency is proved by the former lemma.

Conversely, for each $n \geq 1$, we have the short exact sequence

$$\mathrm{Soc}^n E_x \longrightarrow \mathrm{Soc}^{n+1} E_x \longrightarrow \textstyle\bigoplus_{i \in I_n} S_i \ .$$

If S_y is not a predecessor of S_x then $S_y \ncong S_i$ for all $i \in I_n$. Therefore, for any $n \geq 1$, $\mathrm{Hom}_C(\mathrm{Soc}^{n+1}E_x, E_y) \cong \mathrm{Hom}_C(\mathrm{Soc}^n E_x, E_y)$. Now,

$$\begin{aligned}
\mathrm{Hom}_C(E_x, E_y) &\cong \mathrm{Hom}_C(\varinjlim \mathrm{Soc}^n E_x, E_y) \\
&\cong \varinjlim \mathrm{Hom}_C(\mathrm{Soc}^n E_x, E_y) \\
&\cong \mathrm{Hom}_C(S_x, E_y) \\
&\cong \begin{cases} 0, & \text{if } S_x \ncong S_y, \\ \mathrm{End}_C(S_x), & \text{if } S_x \cong S_y. \end{cases}
\end{aligned}$$

Thus $\mathrm{Rad}_C(E_x, E_y) = 0$. ☐

Theorem 2.9. *Let S_x and S_y be two simple C-comodules and n be a positive integer. If S_y is an n-predecessor of S_x then there exists a path in Γ_C of length n from S_y to S_x.*

Proof. We proceed by induction on the integer n. For $n = 1$ is just Lemma 2.2. Let us assume that the assertion holds for $n - 1$ and let S_y be a n-predecessor of S_x. By Lemma 2.6, there exists a non-zero map $f : \mathrm{Soc}^{n+1}E_x \to E_y$ such that $f(\mathrm{Soc}^n E_x) = 0$. In particular we may decompose f as follows

$$\begin{tikzcd}
\mathrm{Soc}^{n+1}E_x \arrow[r, "f"] \arrow[d, "p"] & E_y \\
M = \frac{\mathrm{Soc}^{n+1}E_x}{\mathrm{Soc}^{n-1}E_x} \arrow[ur, "g"'] &
\end{tikzcd}$$

where g is a non-zero map such that $g(\mathrm{Soc}^n E_x/\mathrm{Soc}^{n-1}E_x) = 0$.

By Lemma 2.4, $M = \mathrm{Soc}^2(E_x/\mathrm{Soc}^{n-1}E_x)$ and then it is contained in $\mathrm{Soc}^2(\bigoplus_{i\in I_{n-1}} E_i) = \bigoplus_{i\in I_{n-1}} \mathrm{Soc}^2 E_i$. Therefore, since E_y is injective, there exists a non-zero map h making commutative the following diagram

$$\begin{tikzcd}
E_y & \\
M \arrow[u, "g"] \arrow[r, "i"] & \bigoplus_{i\in I_{n-1}} \mathrm{Soc}^2 E_i \arrow[ul, "h"']
\end{tikzcd}$$

Hence there is an index $j \in I_{n-1}$ such that the composition $hu_j : \mathrm{Soc}^2 E_j \to E_y$ is non-zero, where u_j is the usual inclusion. Lastly, since $g(\mathrm{Soc}\, M) = 0$, $h(\bigoplus_{i\in I_{n-1}} S_i) = 0$ and then $hu_j(S_j) = 0$. Thus there is an arrow $S_y \to S_j$ in Γ_C. ☐

Remark 2.10. *The reader should observe that if there is a path in Γ_C from S_y to S_x, then S_y does not have to be a predecessor of S_x. For example, consider the quiver Q*

$$\begin{tikzcd}
\underset{y}{\circ} \arrow[r, "\alpha"] & \underset{z}{\circ} \arrow[r, "\beta"] & \underset{x}{\circ}
\end{tikzcd},$$

and the subcoalgebra C of KQ generated by $\{x, y, z, \alpha, \beta\}$. Then the quiver Γ_C is

$$S_y \longrightarrow S_z \longrightarrow S_x.$$

Obviously, there is a path from S_y to S_x, but there is no non-zero morphisms

$$f : E_x =< x, \beta > \longrightarrow E_y =< y > .$$

On the other hand, if C is the coalgebra KQ, the Ext-quiver of KQ is also the previous quiver but, in this case, we may obtain a map

$$f : E_x =< x, \beta, \beta\alpha > \longrightarrow E_y =< y >$$

defined by $f(\beta\alpha) = y$ and zero otherwise.

Lemma 2.11. *Let E_x and E_y be two indecomposable injective right C-comodules and $f : E_x \to E_y$ be a morphism of C-comodules such that $f(\mathrm{Soc}^n E_x) = 0$ and $f(\mathrm{Soc}^{n+1}E_x) \neq 0$. Then $f(\mathrm{Soc}^{n+1}E_x) = S_y$ and $f(\mathrm{Soc}^{n+t}E_x) \subseteq \mathrm{Soc}^t E_y$ for any $t > 1$. Moreover, if C is hereditary, then $f(\mathrm{Soc}^{n+t}E_x) = \mathrm{Soc}^t E_y$ for any $t \geq 1$.*

Proof. We may factorize f as the composition:

$$\begin{tikzcd}
E_x \arrow[r, "p"] \arrow[rr, bend right, "f"'] & \dfrac{E_x}{\mathrm{Soc}^n E_x} \arrow[r, "g"] & E_y
\end{tikzcd}$$

Then $f(\mathrm{Soc}^{n+1}E_x) = gp(\mathrm{Soc}^{n+1}E_x) = g(\frac{\mathrm{Soc}^{n+1}E_x}{\mathrm{Soc}^{n}E_x})$ is a non-zero semisimple subcomodule of E_y, i.e., it is S_y. Let us consider the chain

$$0 \subseteq f(\mathrm{Soc}^n E_x) \subseteq f(\mathrm{Soc}^{n+1}E_x) \subseteq \cdots \subseteq f(\mathrm{Soc}^{n+t}E_x) \subseteq \cdots$$

Since each quotient $\frac{f(\mathrm{Soc}^{n+i}E_x)}{f(\mathrm{Soc}^{n+i-1}E_x)}$ is semisimple for any $i \geq 0$, by Lemma 2.3, $f(\mathrm{Soc}^{n+t}E_x) \subseteq \mathrm{Soc}^t E_y$ for any $t > 1$.

Suppose now that C is hereditary. Then, for each $i \in I_n$, the following diagram commutes

$$\begin{tikzcd}
E_x \arrow[rr, "f"] \arrow[dr, "p"'] & & E_y & \\
& \frac{E_x}{\mathrm{Soc}^n E_x} \cong & \bigoplus_{i\in I_n} E_i \arrow[u, "g"] & E_i \arrow[l, "u_i"] \arrow[ul, "gu_i"']
\end{tikzcd}$$

Since $g(\bigoplus_{i\in I_n} S_i) = S_y$, there is an index $j \in I_n$ such that $g_{|S_j} : S_j \to S_y$ is bijective. Thus gu_j is an isomorphism and

$$\mathrm{Soc}^t E_y = gu_j(\mathrm{Soc}^t E_j) \subseteq g(\bigoplus_{i\in I_n} \mathrm{Soc}^t E_i) = f(\mathrm{Soc}^{n+t}E_x)$$

for any $t > 0$. □

Corollary 2.12. *Let C be a hereditary coalgebra and n be a positive integer. The following conditions are equivalent:*

- (a) *There is a path in Γ_C of length n from a vertex S_y to a vertex S_x.*
- (b) *S_y is an n-predecessor of S_x.*

Proof. It is enough to prove $(a) \Rightarrow (b)$. If

$$S_y \longrightarrow S_1 \longrightarrow \cdots \longrightarrow S_{n-1} \longrightarrow S_x$$

is a path in Γ_C, there exists a sequence of (surjective) morphisms

$$E_x \xrightarrow{f_n} E_{n-1} \xrightarrow{f_{n-1}} \cdots \xrightarrow{f_2} E_1 \xrightarrow{f_1} E_y$$

such that $f_i(S_i) = 0$ and $f_i(\mathrm{Soc}^2 E_i) \neq 0$ for all $i = 1, \ldots, n$, where $S_n = S_x$. Then, applying repeatedly the previous lemma, we obtain that $(f_1 f_2 \cdots f_n)(\mathrm{Soc}^n E_x) = 0$ and $(f_1 f_2 \cdots f_n)(\mathrm{Soc}^{n+1}E_x) = S_y \neq 0$. By Lemma 2.6, S_y is an n-predecessor of S_x. □

3 The section functor

From now on we fix an idempotent element $e \in C^*$. We will denote by $\mathcal{T}_e$ the localizing subcategory associated to e and by $\{S_x\}_{x \in I_e \subset I_C}$ the subset of simple comodules of the quotient category. Let us consider the quotient and the section functor associated to $\mathcal{T}_e$:

$$\begin{tikzcd}
\mathcal{M}^C \arrow[r, shift left, "T=e(-)=-\square_C eC"] & \mathcal{M}^{eCe} \arrow[l, shift left, "S=-\square_{eCe}Ce"]
\end{tikzcd}.$$

We recall that there exists a torsion theory on $\mathcal{M}^C$ associated to the functor T, where a right C-comodule M is a torsion comodule if $T(M) = 0$. If M is not torsion, we denote by $t(M)$ the torsion subcomodule of M.

We know that, by [10], for a simple right C-comodule S_x, $T(S_x) = S_x$ if $x \in I_e$ and zero otherwise. From this fact we obtain the following result:

Lemma 3.1. *Let M be a right C-comodule then $T(\operatorname{Soc} M) \subseteq \operatorname{Soc} T(M)$.*

Proof. Let us suppose that $\operatorname{Soc} M = (\bigoplus_{i\in I} S_i) \bigoplus (\bigoplus_{j \in J} T_j)$, where S_i and T_j are simple right C-comodules such that $T(S_i) = S_i$ and $T(T_j) = 0$ for all $i \in I$ and $j \in J$. Since $\operatorname{Soc} M \subseteq M$ then we have that $T(\operatorname{Soc} M) = \bigoplus_{i \in I} S_i \subseteq T(M)$. $\square$

Let us study the behavior of the injective comodules under the action of the section functor. Indeed, we shall prove that S preserves indecomposable injective comodules and, consequently, injective envelopes. In what follows we will denote by $\{\overline{E}_x\}_{x \in I_e}$ a complete set of pairwise non-isomorphic indecomposable injective right eCe-comodules, and assume that $\overline{E}_x$ is the injective envelope of the simple right eCe-comodule S_x for each $x \in I_e$.

Proposition 3.2. *The following properties hold:*

(*a*) *The functor S preserves injective comodules.*

(*b*) *If N is a quasi-finite indecomposable right eCe-comodule then $S(N)$ is indecomposable.*

(*c*) *The functor S preserves indecomposable injective comodules.*

(*d*) *If S_x is a simple eCe-comodule then* $\operatorname{Soc} S(S_x) = S_x$.

(*e*) *If S_x is a simple eCe-comodule then $S(S_x)$ is torsion-free.*

(*f*) *We have that $S(\overline{E}_x) = E_x$ for all $x \in I_e$.*

(*g*) *The functor S preserves quasi-finite comodules.*

(h) *The functor $S : \mathcal{M}^{eCe} \to \mathcal{M}^{C}$ restricts to a fully faithful functor $S : \mathcal{M}^{eCe}_{qf} \to \mathcal{M}^{C}_{qf}$ between the categories of quasi-finite comodules which preserves indecomposables comodules and respects isomorphism classes.*

Proof. (a) The functor T is exact and left adjoint of S so, by [18, Proposition 9.5], the result follows.

(b) Since N is quasi-finite and indecomposable then we have that $\operatorname{End}_{eCe}(N) \cong \operatorname{End}_{C}(S(N))$ is a local ring. Thus $S(N)$ is indecomposable.

(c) It follows from (a) and (b).

(d) Suppose that $\operatorname{Soc} S(S_x) = (\bigoplus_{i\in I} S_i) \bigoplus (\bigoplus_{j \in J} T_j)$, where S_i and T_j are simple right C-comodules such that $T(S_i) = S_i$ and $T(T_j) = 0$ for all $i \in I$ and $j \in J$. By Lemma 3.1, $\bigoplus_{i \in I} S_i = T(\operatorname{Soc} S(S_x)) \subseteq \operatorname{Soc} TS(S_x) = \operatorname{Soc} S_x = S_x$. Since S is left exact and preserves indecomposable injective comodules, $S_x \subseteq \operatorname{Soc} S(S_x) \subseteq \operatorname{Soc} S(\overline{E}_x) = S_y$ for some simple comodule S_y. Then $S_y = S_x = \operatorname{Soc} S(S_x)$.

(e) If $M \subseteq S(S_x)$ is a non-zero torsion subcomodule of $S(S_x)$ then there exists a simple C-comodule R contained in M such that $T(R) = 0$. But $\operatorname{Soc} S(S_x) = S_x$, so $S_x = R$ and we get a contradiction.

(f) It is easy to see from (c) and (d).

(g) Let M be a quasi-finite right eCe-comodule. The injective envelope of M is a quasi-finite injective comodule $M \hookrightarrow E = \bigoplus \overline{E}_x^{n_x}$. Since S is left exact then $S(M) \hookrightarrow S(E) = \bigoplus E_x^{n_x}$. Thus $S(M)$ is quasi-finite.

(h) It is a consequence of the above assertions and the equality $TS = 1_{\mathcal{M}^{eCe}}$.

□

Corollary 3.3. *S preserves injective envelopes.*

After proving Proposition 3.2, one should ask if the behavior of simple comodules is analogous to injective ones, that is, if S preserves simple comodules and, consequently, in view of Proposition 3.2(c), $S(S_x) = S_x$ for all $x \in I_e$. Unfortunately, in general, this is not true and we can only say that $S(S_x)$ is a subcomodule of E_x which contains S_x.

Example 3.4. *This example shows that $S(S_x)$ does not have to be S_x for every $x \in I_e$. Consider the quiver Q*



$C = KQ$ and the idempotent $e \in C^*$ associated to the set $\{x\}$. Then, the localized coalgebra eCe is S_x and

$$S(S_x) = S_x \square_{eCe} Ce = eCe \square_{eCe} Ce \cong Ce \cong < x, \alpha > \neq S_x.$$

The reader should observe that $S(S_x)$ could be an infinite dimensional right C-comodule. Therefore, in general, S cannot be restricted to a functor between the categories of finite dimensional comodules.

Example 3.5. *Consider the quiver Q*



$C = KQ$ and the idempotent $e \in C^*$ associated to the set $\{1\}$. Then the localized coalgebra eCe is S_1 and

$$S(S_1) = S_1 \square_{eCe} Ce = eCe \square_{eCe} Ce \cong Ce \cong < 1, \{\alpha_1 \cdots \alpha_{n-1}\alpha_n\}_{n \geq 1} > .$$

Remark 3.6. *The reader may find in [9] a proof of the following fact: S preserves finite dimensional comodules if and only $S(S_x)$ is finite dimensional for each $x \in I_e$.*

In order to characterize the simple comodules invariant under the functor S we need the following result. It asserts that the torsion immediate predecessors of a torsion-free vertex S_x in Γ_C are the simple C-comodules contained in the socle of $S(S_x)/S_x$. In the following picture the torsion-free vertices are represented by white points.



Theorem 3.7. *Let S_y and S_x be two simple C-comodules. Then we have that $S_y \subseteq S(S_x)/S_x$ if and only if $S_y \subseteq E_x/S_x$ and $T(S_y) = 0$.*

Proof. Consider the short exact sequence

$$\begin{tikzcd} S_x \arrow[r] & S(S_x) \arrow[r] & S(S_x)/S_x \end{tikzcd} \tag{1}$$

Since $S_x = T(S_x) = TS(S_x)$, $S(S_x)/S_x$ is a torsion subcomodule of E_x/S_x. Therefore if $S_y \subseteq S(S_x)/S_x$ then $S_y \subseteq E_x/S_x$ and $T(S_y) = 0$.

Conversely, applying the functor S to the exact sequence

$$S_x \xrightarrow{i} \overline{E}_x \xrightarrow{p} \overline{E}_x/S_x$$

we obtain the following commutative diagram:

$$\begin{tikzcd}
S(S_x) \arrow[r, "S(i)"] & E_x \arrow[rr, "S(p)"] \arrow[dr, "\mathrm{Coker}\ S(i)"'] & & S(\overline{E}_x/S_x) \arrow[r] & \mathrm{Coker} \\
& & E_x/S(S_x) \arrow[r] \arrow[ur, hook] & 0 &
\end{tikzcd}$$

Therefore we have that $\mathrm{Hom}_C(S_y, E_x/S(S_x))$ is contained in the set of morphisms $\mathrm{Hom}_C(S_y, S(\overline{E}_x/S_x)) \cong \mathrm{Hom}_{eCe}(T(S_y), \overline{E}_x/S_x) = 0$. Now, applying $\mathrm{Hom}_C(S_y, -)$ to the exact sequence

$$S(S_x) \longrightarrow E_x \longrightarrow E_x/S(S_x),$$

we obtain the exactness of the sequence

$$0 = \mathrm{Hom}_C(S_y, E_x) \to \mathrm{Hom}_C(S_y, E_x/S(S_x)) \to \mathrm{Ext}^1_C(S_y, S(S_x)) \to 0$$

and then $0 = \mathrm{Hom}_C(S_y, E_x/S(S_x)) \cong \mathrm{Ext}^1_C(S_y, S(S_x))$.

Let us now apply the functor $\mathrm{Hom}_C(S_y, -)$ to (1) and therefore $\mathrm{Hom}_C(S_y, S(S_x)/S_x) \cong \mathrm{Ext}^1_C(S_y, S_x) \neq 0$. Then the result follows. $\square$

Corollary 3.8. *Let S_x be a simple eCe-comodule. The following conditions are equivalent:*

(*a*) *E_x/S_x is torsion-free.*

(*b*) *There is no arrow in Γ_C from a torsion vertex S_y to S_x.*

(*c*) *$S(S_x) = S_x$.*

Proof. It is straightforward from Lemma 2.2 and Theorem 3.7. $\square$

We may generalize the former results using the Loewy series of the C-comodule $S(S_x)$.

Lemma 3.9. *Let S_x and S_y be two simple C-comodules such that S_x is torsion-free. Then $S_y \subseteq S(S_x)/\mathrm{Soc}^n S(S_x)$ if and only if S_y is torsion and $S_y \subseteq E_x/\mathrm{Soc}^n S(S_x)$.*

Proof. The necessity can be proved as the former theorem. Conversely, since S_y is torsion, by the proof of Theorem 3.7, we have that $\operatorname{Ext}^1_C(S_y, S(S_x)) = 0$. Then, applying the functor $\operatorname{Hom}_C(S_y, -)$ to the short exact sequences

$$\begin{tikzcd}\operatorname{Soc}^n S(S_x) \arrow[r] & E_x \arrow[r] & E_x/\operatorname{Soc}^n S(S_x)\end{tikzcd}$$

$$\begin{tikzcd}\operatorname{Soc}^n S(S_x) \arrow[r] & S(S_x) \arrow[r] & S(S_x)/\operatorname{Soc}^n S(S_x)\end{tikzcd}$$

we obtain that $\operatorname{Hom}_C(S_y, \frac{S(S_x)}{\operatorname{Soc}^n S(S_x)}) \cong \operatorname{Hom}_C(S_y, \frac{E_x}{\operatorname{Soc}^n S(S_x)})$. □

Theorem 3.10. *Let E_x be an indecomposable injective C-comodule such that S_x is torsion-free. If $S_y \subseteq S(S_x)/\operatorname{Soc}^n S(S_x)$ for some $n \geq 1$, then the following assertions hold:*

- (*a*) *S_y is torsion.*
- (*b*) *S_y is a n-predecessor of S_x.*
- (*c*) *There exists a path in Γ_C*

$$\begin{tikzcd}S_y \arrow[r] & S_{n-1} \arrow[r] & \cdots \arrow[r] & S_2 \arrow[r] & S_1 \arrow[r] & S_x\end{tikzcd}$$

such that S_i is torsion for all $i = 1, \ldots, n-1$.

The converse also holds if C is hereditary.

Proof. (*a*) can be proved as in Theorem 3.7. (*b*) is obtained from the inclusion $\frac{S(S_x)}{\operatorname{Soc}^n S(S_x)} \subseteq \frac{E_x}{\operatorname{Soc}^n E_x}$. For (*c*), we proceed by induction on the number n. The case $n = 1$ corresponds to Theorem 3.7. Assume now that the statement holds for $n-1$ and that $S_y \subseteq S(S_x)/\operatorname{Soc}^n S(S_x)$. Analogously to the proof of Theorem 2.9, we may prove that there exists an arrow from S_y to a simple C-comodule $S_j \subseteq S(S_x)/\operatorname{Soc}^{n-1} S(S_x)$.

Let us now suppose that C is hereditary. We will prove that, for each positive integer n, the torsion simple comodules contained in $E_x/\operatorname{Soc}^n S(S_x)$ are those for which there is a path as described in (*c*). By Lemma 3.9, this will imply the statement. The case $n = 1$ is just Theorem 3.7. Let us assume that it is verified for $n-1$. Then

$$\frac{E_x}{\operatorname{Soc}^n S(S_x)} \cong \frac{\frac{E_x}{\operatorname{Soc}^{n-1} S(S_x)}}{\frac{\operatorname{Soc}^n S(S_x)}{\operatorname{Soc}^{n-1} S(S_x)}} \cong \frac{(\oplus_{j\in J} E_j) \oplus \oplus E}{\oplus_{j\in J} S_j} \cong (\oplus_{j\in J} E_j/S_j) \oplus E,$$

where S_j is torsion for all $j \in J$ and E is an injective comodule whose socle is torsion-free. If there is a path from S_y to S_x as described in (*c*), $1 \in J$ by hypothesis and hence $S_y \subseteq E_1/S_1 \subseteq E_x/\operatorname{Soc}^n S(S_x)$. Conversely, there is some $j_o \in J$ such that $S_y \subseteq E_{j_0}/S_{j_0}$ and then we have an arrow $S_y \to S_{j_0}$. By hypothesis there is a path of length $n-1$ as described in (*c*) from S_{j_0} to S_x. This completes the proof. □

Corollary 3.11. *Let Q be a quiver, KQ the path coalgebra of Q and $e \in (KQ)^*$ an idempotent element associated to the subset $X \subseteq Q_0$. For each vertex $x \in X$, the KQ-comodule $S(S_x)$ is generated by the set of paths*

$$\begin{tikzcd} \underset{x_1}{\circ} \arrow[r] & \underset{x_2}{\circ} \arrow[r] & \circ \arrow[r, dash] & \cdots \arrow[r, dash] & \circ \arrow[r] & \underset{x_n}{\circ} \arrow[r] & \underset{x}{\circ} \end{tikzcd}$$

such that $x_i \notin X$ for any $i = 1, \dots, n$

Corollary 3.12. *Suppose that $\dim_K \mathrm{Ext}^1_C(S_1, S_2)$ is finite for each simple C-comodules S_1 and S_2. If there is finitely many paths in Γ_C,*

$$\begin{tikzcd} S_t \arrow[r] & S_{t-1} \arrow[r] & \cdots \arrow[r] & S_2 \arrow[r] & S_1 \arrow[r] & S_x \end{tikzcd}$$

such that S_i is a torsion simple C-comodule for all $i = 1, \dots, t$, then $S(S_x)$ is finite dimensional. If C is hereditary, the converse holds.

Remark 3.13. *If we consider the non-valued Gabriel quiver of C, it is not needed to assume that the groups of extensions between simple comodules have finite dimension.*

4 The quotient functor

Let us now analyze the properties of the quotient functor. We start with an example which shows that, in general, T does not preserve injective comodules.

Example 4.1. *Let Q be the quiver*

$$\begin{tikzcd} \underset{x}{\circ} \arrow[r, "\alpha"] & \underset{y}{\circ} \arrow[r, "\beta"] & \underset{z}{\circ} \end{tikzcd} ,$$

C be the subcoalgebra of KQ generated by $\{x, y, z, \alpha, \beta\}$ and $I_e = \{x, y\}$. The injective right C-comodule E_z is generated by $< z, \beta >$ and $T(E_z) =< \beta >\cong S_y \neq E_y$.

Proposition 4.2. *The following statements hold:*

- *(a)* *$T(E_x) = \overline{E}_x$ for any $x \in I_e$.*
- *(b)* *If E is an injective torsion-free right C-comodule then $T(E)$ is an injective right eCe-comodule.*
- *(c)* *If M is a torsion-free right C-comodule then $\mathrm{Soc}\, M = \mathrm{Soc}\, T(M) = T(\mathrm{Soc}\, M)$.*
- *(d)* *The functor $T : \mathcal{M}^C \to \mathcal{M}^{eCe}$ restricts to a functor $T : \mathcal{M}^C_{qf} \to \mathcal{M}^{eCe}_{qf}$ and a functor $T : \mathcal{M}^C_f \to \mathcal{M}^{eCe}_f$ between the categories of quasi-finite and finite dimensional comodules, respectively.*

Proof. (*a*) By Proposition 3.2, $E_x = S(\overline{E}_x)$ for any $x \in I_e$. Then $T(E_x) = TS(\overline{E}_x) = \overline{E}_x$.

(*b*) It follows from (*a*).

(*c*) Consider the chain $\bigoplus_{x\in I} S_x = \text{Soc } M \subseteq M \subseteq E(M) = \bigoplus_{x\in I} E_x$. Since M is torsion-free then $I \subseteq I_e$. Therefore $\text{Soc } M = \bigoplus_{x\in I} S_x = \bigoplus_{x\in I} T(S_x) = T(\text{Soc } M) \subseteq T(M) \subseteq T(E(M)) = \bigoplus_{x\in I} T(E_x) = \bigoplus_{x\in I} \overline{E}_x$ and the result follows.

(*d*) It is easy to see.

□

Corollary 4.3. *Let E_x be a indecomposable injective C-comodule such that S_x is torsion-free. $S(S_x) = E_x$ if and only if all predecessors of S_x in Γ_C are torsion.*

Proof. Assume that all predecessors of S_x are torsion. Then,

$$T(\text{Soc}^n E_x) = T(\text{Soc}^{n+1} E_x) = T(\text{Soc } E_x) = S_x.$$

Now, by the previous proposition,

$$\overline{E}_x = T(E_x) = T(\varinjlim \text{Soc}^n E_x) = \varinjlim T(\text{Soc}^n E_x) = S_x$$

and thus, by Proposition 3.2(*f*), $E_x = S(\overline{E_x}) = S(S_x)$. The converse follows from Theorem 3.10. □

Example 4.4. *In general, the functor T is not full. Let Q be the quiver*



$C = KQ$ and $e \in C^*$ be the idempotent associated to the set $\{x\}$. Then $\dim_K \text{Hom}_C(S_x, C) = \dim_K \text{End}(S_x) = 1$ *and* $\dim_K \text{Hom}_{eCe}(S_x, eC) = 2$. *Therefore the map* $T_{S_x,C} : \text{Hom}_C(S_x, C) \to \text{Hom}_{eCe}(S_x, eC)$ *cannot be surjective.*

Example 4.5. *In general, the functor T does not preserve indecomposable comodules. Let KQ be the path coalgebra of the quiver*



and $e \in C^*$ be the idempotent associated to the set $\{x, y\}$. Then T maps the indecomposable injective right C-comodule $E_z = < z, \alpha, \beta >$ to the right eCe-comodule $S_x \oplus S_y$. Nevertheless, it is easy to see that T preserves indecomposable torsion-free comodules.

Since $T(S_y) = 0$ for each torsion simple C-comodule, one could expect the analogous property for their injective envelopes. Unfortunately, this is only true in some special conditions related to the stability of the the torsion theory.

Example 4.6. *Let KQ be the path coalgebra of the quiver*



$C = KQ$ and $e \in C^*$ be the idempotent associated to the set $\{x\}$. Then $T(E_y) = T(< y, \alpha >) \cong S_x \neq 0$.

Theorem 4.7. *Let E_y be an indecomposable injective right C-comodule with $y \notin I_e$. The following statements are equivalent:*

- *(a) $T(E_y) = 0$,*
- *(b) $\mathrm{Hom}_C(E_y, E_x) = 0$ for all $x \in I_e$,*
- *(c) S_y has no torsion-free predecessor in Γ_C.*

Proof. $(a) \Rightarrow (b)$. Since S is left adjoint to T then we have that $\mathrm{Hom}_C(E_y, E_x) = \mathrm{Hom}_C(E_y, S(\overline{E}_x)) \cong \mathrm{Hom}_{eCe}(T(E_y), \overline{E}_x) = 0$ for all $x \in I_e$.

$(b) \Rightarrow (c)$. It is proved in Proposition 2.8.

$(c) \Rightarrow (a)$. For each $n \geq 1$, we have the short exact sequence

$$\mathrm{Soc}^n E_y \longrightarrow \mathrm{Soc}^{n+1} E_y \longrightarrow \textstyle\bigoplus_{i \in I_n} S_i \ .$$

Since S_y has no torsion-free predecessor, $T(S_i) = 0$ for all $i \in I_n$ and then $T(\mathrm{Soc}^n E_y) = T(\mathrm{Soc}^{n+1} E_y)$. Now,

$$T(E_y) = T(\varinjlim \mathrm{Soc}^n E_y) = \varinjlim T(\mathrm{Soc}^n E_y) = T(\mathrm{Soc}\ E_y) = 0.$$

□

Let us finish the section by giving an approach to the image of an indecomposable injective comodule E_y with torsion socle. Firstly, from the Loewy series of E_y, we may obtain a chain

$$0 \subseteq T(\mathrm{Soc}E_y) \subseteq T(\mathrm{Soc}^2 E_y) \subseteq \cdots \subseteq T(\mathrm{Soc}^n E_y) \subseteq \cdots \subseteq T(E_y)$$

such that each quotient $\frac{T(\mathrm{Soc}^{n+1} E_y)}{T(\mathrm{Soc}^n E_y)} \cong T\left(\frac{\mathrm{Soc}^{n+1} E_y}{\mathrm{Soc}^n E_y}\right)$ is the the direct sum of the torsion-free n-predecessors of S_y. As a consequent, by Lemma 2.3, we have that $T(\mathrm{Soc}^{n+1} E_y) \subseteq \mathrm{Soc}^n T(E_y)$. In particular, $T(\mathrm{Soc}^2 S_y)$ is the direct sum of all torsion-free immediate predecessors of S_y and $T(\mathrm{Soc}^2 E_y) \subseteq \mathrm{Soc}\, T(E_y)$.

Lemma 4.8. *Let S_y be a torsion simple right C-comodule. Suppose that $\{S_x, T_z\}_{x\in I, z\in J}$ is the set of all immediate predecessors of S_y in Γ_C, where S_x is torsion-free for all $x \in I$ and T_z is torsion for all $z \in J$. Then*

$$\operatorname{Soc} T(E_y) \subseteq \left(\bigoplus_{x\in I} S_x\right) \bigoplus \left(\bigoplus_{z\in J} \operatorname{Soc} T(E_z)\right).$$

If C is hereditary, the opposite inclusion also holds.

Proof. By Lemma 2.2, $\operatorname{Soc}(E_y/S_y) = (\oplus_{x\in I} S_x) \oplus (\oplus_{z\in J} T_z)$ and, consequently, $E_y/S_y \subseteq (\oplus_{x\in I} E_x) \oplus (\oplus_{z\in J} E_z)$. Then

$$T(E_y) \cong T(E_y/S_y) \subseteq \left(\bigoplus_{x\in I} \overline{E}_x\right) \bigoplus \left(\bigoplus_{z\in J} T(E_z)\right)$$

and then

$$\operatorname{Soc} T(E_y) \subseteq \left(\bigoplus_{x\in I} S_x\right) \bigoplus \left(\bigoplus_{z\in J} \operatorname{Soc} T(E_z)\right).$$

Clearly, the inclusions are equalities if C is hereditary. ☐

In a general context it is not possible to prove the equality in Lemma 4.8. For example, consider the quiver of Example 4.1, the coalgebra generated by the set $< x, y, z, \alpha, \beta >$ and $I_e = \{x\}$. Then $\operatorname{Soc} T(E_z) = 0$ and $\operatorname{Soc} T(E_y) = S_x$.

Corollary 4.9. *Let E_y be a indecomposable injective C comodule such that S_y is torsion. If $S_x \subseteq \operatorname{Soc} T(E_y)$ then*

- (a) *S_x is torsion-free.*
- (b) *S_x is a predecessor of S_y in Γ_C.*
- (c) *There exists a path in Γ_C*

$$\begin{tikzcd} S_x \arrow[r] & S_n \arrow[r] & \cdots \arrow[r] & S_2 \arrow[r] & S_1 \arrow[r] & S_y \end{tikzcd}$$

such that S_i is torsion for all $i = 1, \dots, n$.

If C is hereditary, the converse also holds.

Proof. By Lemma 4.8, it is enough to prove (b). Now, by hypothesis, $0 \neq \operatorname{Hom}_{eCe}(T(E_y), \overline{E}_x) \cong \operatorname{Hom}_C(E_y, E_x) = \operatorname{Rad}_C(E_y, E_x)$. Corollary 2.8 completes the proof. ☐

5 The colocalizing functor

Throughout this section we shall assume that $\mathcal{T}_e$ is a colocalizing subcategory of $\mathcal{M}^C$. Then the quotient functor T has a left adjoint functor H which can be described as follows:

$$\begin{tikzcd}\mathcal{M}^C \arrow[r, shift left, "T=e(-)=-\square_C eC"] & \mathcal{M}^{eCe} \arrow[l, shift left, "H=\mathrm{Cohom}_{eCe}(eC,-)"] \end{tikzcd} .$$

We recall from [20] that there exists such functor H if and only if $eC = T(C)$ is quasi-finite as right eCe-comodule, i.e., since $T(C) = eCe \bigoplus(\bigoplus_{y\in I_C\setminus I_e} T(E_y))$, if and only if $\bigoplus_{y\in I_C\setminus I_e} T(E_y)$ is quasi-finite as eCe-comodule. According to Corollary 4.9, this is obtained if $\dim_K \mathrm{Ext}^1_C(S,S')$ is finite for each pair of simple comodules S and S', and there is finitely many paths in Γ_C

$$\begin{tikzcd} S_x \arrow[r] & S_n \arrow[r] & \cdots \arrow[r] & S_3 \arrow[r] & S_2 \arrow[r] & S_1 \end{tikzcd}$$

where S_i is torsion for all $i = 1,\ldots,n$, for each torsion-free simple comodule S_x.

Proposition 5.1. *The following assertions hold:*

- (a) *H preserves projective comodules.*
- (b) *H preserves finite dimensional comodules.*
- (c) *H preserves finite dimensional indecomposable comodules.*
- (d) *The functor $H : \mathcal{M}^{eCe} \to \mathcal{M}^C$ restricts to a fully faithful functor $H : \mathcal{M}^{eCe}_f \to \mathcal{M}^C_f$ between the categories of finite-dimensional comodules which preserves indecomposable comodules and respects isomorphism classes.*

Proof. (a) It is symmetric to the proof of Proposition 3.2(a).

- (b) Let N be a finite dimensional right eCe-comodule. Then $H(N) = \mathrm{Cohom}_{eCe}(eC,N) = \varinjlim \mathrm{Hom}_{eCe}(N_\lambda, eC)^* = \mathrm{Hom}_{eCe}(N,eC)^*$. Now, since eC is a quasi-finite right eCe-comodule, $\mathrm{Hom}_{eCe}(N,eC)$ has finite dimension.
- (c) Let N be a finite dimensional indecomposable right eCe-comodule. Since H is fully faithful then $\mathrm{End}_{eCe}(N) \cong \mathrm{End}_C(H(N))$ is a local ring. Now, by (b), $H(N)$ is finite dimensional and then $H(N)$ is indecomposable.
- (d) It is straightforward from (b), (c) and the equality $TH = 1_{\mathcal{M}^{eCe}}$.

□

Analogously to the study of the section functor, let us characterize the simple comodules which are invariant under the functor H. For that purpose we need the following proposition:

Proposition 5.2. *Let S_x be a simple eCe-comodule. Then $H(S_x) = S_x$ if and only if* $\mathrm{Hom}_{eCe}(S_x, T(E_y)) = 0$ *for all $y \notin I_e$.*

Proof. From the decomposition $eC = eCe \oplus eC(1-e)$ as right eCe-comodules, we have the following equalities:

$$\begin{aligned}
\dim_K H(S_x) = \; & \dim_K \mathrm{Cohom}_{eCe}(eC, S_x) \\
= \; & \dim_K S_x + \dim_K \mathrm{Cohom}_{eCe}(eC(1-e), S_x) \\
= \; & \dim_K S_x + \textstyle\sum_{y \notin I_e} \dim_K \mathrm{Cohom}_{eCe}(T(E_y), S_x) \\
= \; & \dim_K S_x + \textstyle\sum_{y \notin I_e} \dim_K \mathrm{Hom}_{eCe}(S_x, T(E_y)).
\end{aligned}$$

Therefore, $\mathrm{Hom}_{eCe}(S_x, T(E_y)) = 0$ for all $y \notin I_e$ if $H(S_x) \cong S_x$.

Conversely, there is a natural isomorphism $\mathrm{Hom}_C(H(S_x), S_x) \cong \mathrm{Hom}_{eCe}(S_x, S_x)$ so there exists a non-zero (and then surjective) morphism $f : H(S_x) \to S_x$. By hypothesis, $dim_K H(S_x) = \dim_K S_x$ and then f is an isomorphism. □

Theorem 5.3. *Let S_x be a simple eCe-comodule. $H(S_x) = S_x$ if and only if* $\mathrm{Ext}^1_C(S_x, S_y) = 0$ *for all $y \notin I_e$, i.e., there is no arrow $S_x \to S_y$ in Γ_C, where S_y is a torsion simple C-comodule.*

Proof. By Proposition 5.2, it is enough to prove that $\mathrm{Ext}^1_C(S_x, S_y) = 0$ for all $y \notin I_e$ if and only if $\mathrm{Hom}_{eCe}(S_x, T(E_y)) = 0$ for all $y \notin I_e$.

$\Leftarrow$) Suppose that $\mathrm{Ext}^1_C(S_x, S_y) \neq 0$ for some $y \notin I_e$. By Lemma 2.2, $S_x \subseteq \mathrm{Soc}\,(E_y/S_y)$. Then

$$S_x = T(S_x) \subseteq T(\mathrm{Soc}\, E_y/S_y) \subseteq \mathrm{Soc}\, T(E_y/S_y) = \mathrm{Soc}\, T(E_y)$$

and therefore $\mathrm{Hom}_{eCe}(S_x, T(E_y)) \neq 0$.

$\Rightarrow$) If $\mathrm{Ext}^1_C(S_x, S_y) = 0$ for each $y \notin I_e$ then there is no path as described in Corollary 4.9(c). Thus S_x is not contained in $T(E_y)$ for any $y \notin I_e$.

□

6 Semicentral idempotents

In this section we extend the result obtained in [10] which characterize stable subcategories as those whose associated idempotent is left semicentral. We recall from [10] that an idempotent element $e \in C^*$ is said to be *left (right) semicentral* if $eC = eCe$ ($Ce = eCe$), or equivalently, if eC (Ce) is a subcoalgebra of C. A localizing subcategory is said to be *stable* if it is closed for injective envelopes.

In the following theorem we describe stable subcategories from different points of view. A proof of some equivalences is given in [10, Theorem 4.3]. We recall that, for a subset Λ of the vertex set $(\Gamma_C)_0$, we say that Λ is *right link-closed* if it satisfies that, for each arrow $S \to T$ in Γ_C, if $S \in \Lambda$ then $T \in \Lambda$.

Theorem 6.1. *Let C be a coalgebra and $\mathcal{T}_e \subseteq \mathcal{M}^C$ be a localizing subcategory associated to an idempotent element $e \in C^*$. The following conditions are equivalent:*

- (a) *$\mathcal{T}_e$ is a stable subcategory.*
- (b) *$T(E_x) = 0$ for any $x \notin I_e$.*
- (c) $T(E_x) = \begin{cases} \overline{E}_x & \text{if } x \in I_e, \\ 0 & \text{if } x \notin I_e. \end{cases}$
- (d) *$\mathrm{Hom}_C(E_y, E_x) = 0$ for all $x \in I_e$ and $y \notin I_e$.*
- (e) *Any torsion vertex in Γ_C has no torsion-free predecessor.*
- (f) *$\mathcal{K} = \{S \in (\Gamma_C)_0 \mid T(S) = S\}$ is a right link-closed subset of $(\Gamma_C)_0$, i.e., there is no arrow $S_x \to S_y$ in Γ_C, where $T(S_x) = S_x$ and $T(S_y) = 0$.*
- (g) *There is no path in Γ_C from a vertex S_x to a vertex S_y such that $T(S_x) = S_x$ and $T(S_y) = 0$.*
- (h) *e is a left semicentral idempotent in C^*.*

If T_e is a colocalizing subcategory these are also equivalent to

- (i) *$H(S_x) = S_x$ for any $x \in I_e$.*

Proof. $(a) \Rightarrow (b) \Rightarrow (c) \Rightarrow (d) \Rightarrow (e) \Rightarrow (f) \Rightarrow (g)$. It follows from the definition, Theorem 4.7 and Proposition 2.8.
$(g) \Rightarrow (f)$. Trivial.
$(f) \Rightarrow (c)$. By Corollary 4.9, $\mathrm{Soc}\, T(E_y) = 0$ for all $y \notin I_e$. Then $T(E_y) = 0$ for all $y \notin I_e$.
$(c) \Rightarrow (a)$. Let M be a torsion right C-comodule such that its injective envelope is $\bigoplus_{i \in J} E_i$. Then $S_i \subseteq M$ is torsion for all $i \in J$ and, by hypothesis, $T(E_i) = 0$ for all $i \in J$. Thus $T(\bigoplus_{i \in J} E_i) = 0$.
$(c) \Leftrightarrow (h)$. We have that $C = \bigoplus_{x \in I_C} E_x$ and therefore $T(C) = (\bigoplus_{x \in I_e} \overline{E}_x) \bigoplus (\bigoplus_{y \notin I_e} T(E_y))$. On the other hand, $eCe = \bigoplus_{x \in I_e} \overline{E}_x$. Therefore if (c) holds then $eCe = T(C)$. Conversely, if $eCe = T(C)$ then $\bigoplus_{x \in I_e} \overline{E}_x = (\bigoplus_{x \in I_e} \overline{E}_x) \bigoplus (\bigoplus_{y \notin I_e} T(E_y))$. Since eCe is quasi-finite, by Krull-Remak-Schmidt-Azumaya theorem, $T(E_y) = 0$ for all $y \notin I_e$.
$(e) \Leftrightarrow (i)$. It is Theorem 5.3. ☐

We may find an analogous result to Theorem 6.1 for right semicentral idempotents:

Theorem 6.2. *Let C be a coalgebra and $\mathcal{T}_e \subseteq \mathcal{M}^C$ be a localizing subcategory associated to an idempotent element $e \in C^*$. The following conditions are equivalent:*

- (a) *$\mathcal{T}_{1-e}$ is a stable subcategory.*
- (b) *$T(E_x) = E_x$ for any $x \in I_e$.*
- (c) *$\mathrm{Hom}_C(E_x, E_y) = 0$ for all $x \in I_e$ and $y \notin I_e$.*
- (d) *Any torsion-free vertex in Γ_C has no torsion predecessor.*
- (e) *There is no path in Γ_C from a vertex S_y to a vertex S_x such that $T(S_x) = S_x$ and $T(S_y) = 0$.*
- (f) *$\mathcal{K} = \{S \in (\Gamma_C)_0 \mid T(S) = S\}$ is a left link-closed subset of $(\Gamma_C)_0$, i.e., there is no arrow $S_y \to S_x$ in Γ_C, where $T(S_x) = S_x$ and $T(S_y) = 0$.*
- (g) *e is a right semicentral idempotent in C^*.*
- (h) *The torsion subcomodule of a right C-comodule M is $(1-e)M$*
- (i) *$S(S_x) = S_x$ for all $x \in I_e$.*

Proof. By [10] and Theorem 6.1, it is easy to prove $(a) \Leftrightarrow (b) \Leftrightarrow (c) \Leftrightarrow (d) \Leftrightarrow (e) \Leftrightarrow (f)$.

$(d) \Leftrightarrow (g)$. It is Corollary 3.8. □

As a consequence, for a non necessarily indecomposable coalgebra C, we get the following immediate result:

Corollary 6.3. *The following are equivalent:*

- (a) *e is a central idempotent.*
- (b) *For each arrow $S_x \to S_y$ in Γ_C, $T(S_x) = 0$ if and only if $T(S_y) = 0$.*
- (c) *For each connected component of Γ_C, either all vertices are torsion or all vertices are torsion-free.*
- (d) *$T(E_x) = 0$ for any $x \notin I_e$ and $S(S_x) = S_x$ for all $x \in I_e$.*

Corollary 6.4. *Let e be a central idempotent in C^* and $\{\Sigma_C^t, \Delta_C^s\}_{t,s}$ be the connected components of Γ_C, where the vertices of each Σ_C^t are torsion-free and the vertices of each Δ_C^s are torsion. Then $\{\Sigma_C^t\}_t$ are the connected components of Γ_{eCe}.*

References

[1] E. Abe, Hopf Algebras, Cambridge University Press, 1977.

[2] J. Cuadra and J. Gómez-Torrecillas, Idempotents and Morita-Takeuchi theory, Comm. Algebra, 30 (2002), no. 5, 2405-2426.

[3] J. Cuadra and J. Gómez-Torrecillas, Serial coalgebras, Journal Pure Appl. Algebra, 189 (2004), 89-107.

[4] S. Dăscălescu, C. Năstăsescu and S. Raianu, Hopf algebras. An introduction. Monographs and Textbooks in Pure and Applied Mathematics, 235. Marcel Dekker, New York, 2001.

[5] P. Gabriel, Des categories abeliennes, Bull. Soc. Math. France, 1962, 90, 323-448.

[6] J. A. Green, Locally finite representations, J. Algebra 41, 137-171 (1976).

[7] P. Jara, D. Llena, L. M. Merino and Dragoş Ştefan, Herederitary and formally smooth coalgebras, Algebr. Represent. Theory 8 (2005), no. 3, 363–374.

[8] P. Jara, L. M. Merino and G. Navarro, On path coalgebras of quivers with relations, Colloq. Math. 102 (2005), 49–65.

[9] P. Jara, L. M. Merino and G. Navarro, Localization of tame and wild coalgebras, preprint, 22 pages, 2006.

[10] P. Jara, L. M. Merino, G. Navarro and F. Ruiz, Localization in coalgebras. Stable localizations and path coalgebras, Comm. Algebra, to appear. 2006.

[11] J. Kosakowska and D. Simson, Hereditary coalgebras and representations of species, J. Algebra, 293 (2005) 457-505.

[12] I.-P. Lin, Semiperfect coalgebras, J. Algebra, 49 (1977), 357-373.

[13] S. Montgomery, Hopf Algebras and Their Actions on Rings, MBS, No. 82, AMS, 1993.

[14] S. Montgomery, Indescomposable coalgebras, simple comodules and pointed Hopf algebras. Proc. A.M.S. 123(1995), 2343–2351.

[15] S. Nowak and D. Simson, Locally Dynkin quivers and hereditary coalgebras whose left comodules are direct sums of finite dimensional comodules, Comm. Algebra 30 (2002), no. 1, 455–476.

[16] C. Năstăsescu and B. Torrecillas, Colocalization on Grothendieck categories with applications to coalgebras, J. Algebra 185 (1996), no. 1, 108–124.

[17] D. Simson, Coalgebras, comodules, pseudocompact algebras and tame comodule type, Colloq. Math., 90 (2001), 101–150.

[18] B. Stenström, Rings of Quotients, Springer-Verlag, 1975.

[19] M. E. Sweedler, Hoft Algebras, Benjamin, New york, 1969.

[20] M. Takeuchi, Morita theorems for categories of comodules, J. Fac. Sci. Uni. Tokyo 24 (1977), 629–644.

[21] D. Woodcock, Some categorical remarks on the representation theory of coalgebras, Comm. Algebra, 25 (1997), 2775–2794.